

A COMBINATORIAL PROOF OF A FORMULA FOR BETTI NUMBERS OF A STACKED POLYTOPE

SUYOUNG CHOI AND JANG SOO KIM

ABSTRACT. For a simplicial complex Δ , the graded Betti number $\beta_{i,j}(\mathbf{k}[\Delta])$ of the Stanley-Reisner ring $\mathbf{k}[\Delta]$ over a field \mathbf{k} has a combinatorial interpretation due to Hochster. Terai and Hibi showed that if Δ is the boundary complex of a d -dimensional stacked polytope with n vertices for $d \geq 3$, then $\beta_{k-1,k}(\mathbf{k}[\Delta]) = (k-1)\binom{n-d}{k}$. We prove this combinatorially.

1. INTRODUCTION

A *simplicial complex* Δ on a finite set V is a collection of subsets of V satisfying

- (1) if $v \in V$ then $\{v\} \in \Delta$,
- (2) if $F \in \Delta$ and $F' \subset F$, then $F' \in \Delta$.

Each element $F \in \Delta$ is called a *face* of Δ . The *dimension* of F is defined by $\dim(F) = |F| - 1$. The *dimension* of Δ is defined by $\dim(\Delta) = \max\{\dim(F) : F \in \Delta\}$. For a subset $W \subset V$, let Δ_W denote the simplicial complex $\{F \cap W : F \in \Delta\}$ on W .

Let Δ be a simplicial complex on V . Two elements $v, u \in V$ are said to be *connected* if there is a sequence of vertices $v = u_0, u_1, \dots, u_r = u$ such that $\{u_i, u_{i+1}\} \in \Delta$ for all $i = 0, 1, \dots, r-1$. A *connected component* C of Δ is a maximal nonempty subset of V such that every two elements of C are connected.

Let $V = \{x_1, x_2, \dots, x_n\}$ and let R be the polynomial ring $\mathbf{k}[x_1, \dots, x_n]$ over a fixed field \mathbf{k} . Then R is a graded ring with the standard grading $R = \bigoplus_{i \geq 0} R_i$. Let $R(-j) = \bigoplus_{i \geq 0} (R(-j))_i$ be the graded module over R with $(R(-j))_i = R_{j+i}$. The *Stanley-Reisner ring* $\mathbf{k}[\Delta]$ of Δ over \mathbf{k} is defined to be R/I_Δ , where I_Δ is the ideal of R generated by the monomials $x_{i_1}x_{i_2} \cdots x_{i_r}$ such that $\{x_{i_1}, x_{i_2}, \dots, x_{i_r}\} \notin \Delta$. A *finite free resolution* of $\mathbf{k}[\Delta]$ is an exact sequence

$$(1) \quad 0 \longrightarrow F_r \xrightarrow{\phi_r} F_{r-1} \xrightarrow{\phi_{r-1}} \cdots \xrightarrow{\phi_2} F_1 \xrightarrow{\phi_1} F_0 \xrightarrow{\phi_0} \mathbf{k}[\Delta] \longrightarrow 0,$$

where $F_i = \bigoplus_{j \geq 0} R(-j)^{\beta_{i,j}}$ and each ϕ_i is degree-preserving. A finite free resolution (1) is *minimal* if each $\beta_{i,j}$ is smallest possible. There is a minimal finite free resolution of $\mathbf{k}[\Delta]$ and it is unique up to isomorphism. If (1) is minimal, then the (i, j) -th *graded Betti number* $\beta_{i,j}(\mathbf{k}[\Delta])$ of $\mathbf{k}[\Delta]$ is defined to be $\beta_{i,j}(\mathbf{k}[\Delta]) = \beta_{i,j}$. Hochster's theorem says

$$\beta_{i,j}(\mathbf{k}[\Delta]) = \sum_{\substack{W \subset V \\ |W|=j}} \dim_{\mathbf{k}} \tilde{H}_{j-i-1}(\Delta_W; \mathbf{k}).$$

We refer the reader to [1, 5] for the details of Betti numbers and Hochster's theorem. Since $\dim_{\mathbf{k}} \tilde{H}_0(\Delta_W; \mathbf{k})$ is the number of connected components of Δ_W minus 1, we can interpret $\beta_{i-1,i}(\mathbf{k}[\Delta])$ in a purely combinatorial way.

Definition 1.1. Let Δ be a simplicial complex on a finite nonempty set V . Let k be a nonnegative integer. The k -th special graded Betti number $b_k(\Delta)$ of Δ is defined to be

$$(2) \quad b_k(\Delta) = \sum_{\substack{W \subset V \\ |W|=k}} (\text{cc}(\Delta_W) - 1),$$

where $\text{cc}(\Delta_W)$ denotes the number of connected components of Δ_W .

Note that since there is no connected component in $\Delta_\emptyset = \{\emptyset\}$, we have $b_0(\Delta) = -1$. If $k > |V|$, then $b_k(\Delta) = 0$ because there is nothing in the sum in (2). Thus we have

$$b_k(\Delta) = \begin{cases} \beta_{k-1,k}(\mathbf{k}[\Delta]), & \text{if } k \geq 1, \\ -1, & \text{if } k = 0. \end{cases}$$

We refer the reader to [7] for the basic notions of convex polytopes. Let P be a simplicial polytope with vertex set V . The *boundary complex* $\Delta(P)$ is the simplicial complex Δ on V such that $F \in \Delta$ for some $F \subset V$ if and only if $F \neq V$ and the convex hull of F is a face of P . Note that if the dimension of P is d , then $\dim(\Delta(P)) = d - 1$.

For a d -dimensional simplicial polytope P , we can attach a d -dimensional simplex to a facet of P . A *stacked polytope* is a simplicial polytope obtained in this way starting with a d -dimensional simplex.

Let P be a d -dimensional stacked polytope with n vertices. Hibi and Terai [6] showed that $\beta_{i,j}(\mathbf{k}[\Delta(P)]) = 0$ unless $i = j - 1$ or $i = j - d + 1$. Since $\beta_{i-1,i}(\mathbf{k}[\Delta(P)]) = \beta_{n-i-d+1,n-i}(\mathbf{k}[\Delta(P)])$, it is sufficient to determine $\beta_{i-1,i}(\mathbf{k}[\Delta(P)])$ to find all $\beta_{i,j}(\mathbf{k}[\Delta(P)])$. In the same paper, they found the following formula for $\beta_{k-1,k}(\mathbf{k}[\Delta(P)])$:

$$(3) \quad \beta_{k-1,k}(\mathbf{k}[\Delta(P)]) = (k - 1) \binom{n - d}{k}.$$

Herzog and Li Marzi [4] gave another proof of (3).

The main purpose of this paper is to prove (3) combinatorially. In the meanwhile, we get as corollaries the results of Bruns and Hibi [2] : a formula of $b_k(\Delta)$ if Δ is a tree (or a cycle) considered as a 1-dimensional simplicial complex.

2. DEFINITION OF t -CONNECTED SUM

In this section we define a t -connected sum of simplicial complexes, which gives another equivalent definition of the boundary complex of a stacked polytope. See [3] for the details of connected sums. And then, we extend the definition of t -connected sum to graphs, which has less restrictions on the construction. Every graph in this paper is simple.

2.1. A t -connected sum of simplicial complexes. Let V and V' be finite sets. A *relabeling* is a bijection $\sigma : V \rightarrow V'$. If Δ is a simplicial complex on V , then $\sigma(\Delta) = \{\sigma(F) : F \in \Delta\}$ is a simplicial complex on V' .

Definition 2.1. Let Δ_1 and Δ_2 be simplicial complexes on V_1 and V_2 respectively. Let $F_1 \in \Delta_1$ and $F_2 \in \Delta_2$ be maximal faces with $|F_1| = |F_2|$. Let V'_2 be a finite set and $\sigma : V_2 \rightarrow V'_2$ a relabeling such that $V_1 \cap V'_2 = F_1$ and $\sigma(F_2) = F_1$. Then the *connected sum* $\Delta_1 \#_{\sigma}^{F_1, F_2} \Delta_2$ of Δ_1 and Δ_2 with respect to (F_1, F_2, σ) is the simplicial complex $(\Delta_1 \cup \sigma(\Delta_2)) \setminus \{F_1\}$ on $V_1 \cup V'_2$. If $\Delta = \Delta_1 \#_{\sigma}^{F_1, F_2} \Delta_2$ and $|F_1| = |F_2| = t$, then we say that Δ is a *t -connected sum* of Δ_1 and Δ_2 .




FIGURE 1. The 1-skeleton of a 2-connected sum of Δ_1 and Δ_2 is not a 2-connected sum of $G(\Delta_1)$ and $G(\Delta_2)$.

Note that if Δ_1 and Δ_2 are $(d-1)$ -dimensional pure simplicial complexes, i.e. the dimension of each maximal face is $d-1$, then we can only define a d -connected sum of them.

Let $\Delta_1, \Delta_2, \dots, \Delta_n$ be simplicial complexes. A simplicial complex Δ is said to be a t -connected sum of $\Delta_1, \Delta_2, \dots, \Delta_n$ if there is a sequence of simplicial complexes $\Delta'_1, \Delta'_2, \dots, \Delta'_n$ such that $\Delta'_1 = \Delta_1$, Δ'_i is a t -connected sum of Δ'_{i-1} and Δ_i for $i = 2, 3, \dots, n$, and $\Delta'_n = \Delta$.

2.2. A t -connected sum of graphs. Let G be a graph with vertex set V and edge set E . Let $W \subset V$. Then the *induced subgraph* $G|_W$ of G with respect to W is the graph with vertex set W and edge set $\{\{x, y\} \in E : x, y \in W\}$. Let

$$b_k(G) = \sum_{\substack{W \subset V \\ |W|=k}} (\text{cc}(G|_W) - 1),$$

where $\text{cc}(G|_W)$ denotes the number of connected components of $G|_W$.

Let Δ be a simplicial complex on V . The 1-skeleton $G(\Delta)$ of Δ is the graph with vertex set V and edge set $E = \{F \in \Delta : |F| = 2\}$. By definition, the connected components of Δ_W and $G(\Delta)|_W$ are identical for all $W \subset V$. Thus $b_k(\Delta) = b_k(G(\Delta))$.

Now we define a t -connected sum of two graphs.

Definition 2.2. Let G_1 and G_2 be graphs with vertex sets V_1 and V_2 , and edge sets E_1 and E_2 respectively. Let $F_1 \subset V_1$ and $F_2 \subset V_2$ be sets of vertices such that $|F_1| = |F_2|$, and $G_1|_{F_1}$ and $G_2|_{F_2}$ are complete graphs. Let V'_2 be a finite set and $\sigma : V_2 \rightarrow V'_2$ a relabeling such that $V_1 \cap V'_2 = F_1$ and $\sigma(F_2) = F_1$. Then the *connected sum* $G_1 \#_{\sigma}^{F_1, F_2} G_2$ of G_1 and G_2 with respect to (F_1, F_2, σ) is the graph with vertex set $V_1 \cup V'_2$ and edge set $E_1 \cup \sigma(E_2)$, where $\sigma(E_2) = \{\{\sigma(x), \sigma(y)\} : \{x, y\} \in E_2\}$. If $G = G_1 \#_{\sigma}^{F_1, F_2} G_2$ and $|F_1| = |F_2| = t$, then we say that G is a t -connected sum of G_1 and G_2 .

Note that in contrary to the definition of t -connected sum of simplicial complexes, it is not required that F_1 and F_2 are maximal, and we do not remove any element in $E_1 \cup \sigma(E_2)$. We define a t -connected sum of G_1, G_2, \dots, G_n as we did for simplicial complexes.

It is easy to see that, if $|F_1| = |F_2| \geq 3$ then $G(\Delta_1 \#_{\sigma}^{F_1, F_2} \Delta_2) = G(\Delta_1) \#_{\sigma}^{F_1, F_2} G(\Delta_2)$. Thus we get the following proposition.

Proposition 2.3. *For $t \geq 3$, if Δ is a t -connected sum of $\Delta_1, \Delta_2, \dots, \Delta_n$, then $G(\Delta)$ is a t -connected sum of $G(\Delta_1), G(\Delta_2), \dots, G(\Delta_n)$.*

Note that Proposition 2.3 is not true if $t = 2$ as the following example shows.

Example 2.4. Let $\Delta_1 = \{12, 23, 13\}$ and $\Delta_2 = \{13, 34, 14\}$ be simplicial complexes on $V_1 = \{1, 2, 3\}$ and $V_2 = \{1, 3, 4\}$. Here 12 means the set $\{1, 2\}$. Let $F_1 = F_2 = \{1, 3\}$ and let σ

be the identity map from V_2 to itself. Then the edge set of $G(\Delta_1 \#_{\sigma}^{F_1, F_2} \Delta_2)$ is $\{12, 23, 34, 14\}$, but the edge set of $G(\Delta_1) \#_{\sigma}^{F_1, F_2} G(\Delta_2)$ is $\{12, 23, 34, 14, 13\}$. See Figure 1.

3. MAIN RESULTS

In this section we find a formula of $b_k(G)$ for a graph G which is a t -connected sum of two graphs. To do this let us introduce the following notation. For a graph G with vertex set V , let

$$c_k(G) = \sum_{\substack{W \subset V \\ |W|=k}} \text{cc}(G|_W).$$

Note that $c_k(G) = b_k(G) + \binom{|V|}{k}$.

Lemma 3.1. *Let G_1 and G_2 be graphs with n_1 and n_2 vertices respectively. Let t be a positive integer and let G be a t -connected sum of G_1 and G_2 . Then*

$$\begin{aligned} c_k(G) = & \sum_{i=0}^k \left(c_i(G_1) \binom{n_2-t}{k-i} + c_i(G_2) \binom{n_1-t}{k-i} \right) \\ & - \binom{n_1+n_2-t}{k} + \binom{n_1+n_2-2t}{k}. \end{aligned}$$

Proof. Let V_1 (resp. V_2) be the vertex set of G_1 (resp. G_2). We have $G = G_1 \#_{\sigma}^{F_1, F_2} G_2$ for some $F_1 \subset V_1$, $F_2 \subset V_2$, a vertex set V'_2 and a relabeling $\sigma : V_1 \rightarrow V'_2$ such that $V_1 \cap V'_2 = F_1$, $\sigma(F_2) = F_1$, and $G_1|_{F_1}$ and $G_2|_{F_2}$ are complete graphs on t vertices.

Let A be the set of pairs (C, W) such that $W \subset V_1 \cup V'_2$, $|W| = k$ and C is a connected component of $G|_W$. Let

$$A_1 = \{(C, W) \in A : C \cap V_1 \neq \emptyset\}, \quad A_2 = \{(C, W) \in A : C \cap V'_2 \neq \emptyset\}.$$

Then $c_k(G) = |A| = |A_1| + |A_2| - |A_1 \cap A_2|$. It is sufficient to show that $|A_1| = \sum_{i=0}^k c_i(G_1) \binom{n_2-t}{k-i}$, $|A_2| = \sum_{i=0}^k c_i(G_2) \binom{n_1-t}{k-i}$ and $|A_1 \cap A_2| = \binom{n_1+n_2-t}{k} - \binom{n_1+n_2-2t}{k}$.

Let B_1 be the set of triples (C_1, W_1, X) such that $W_1 \subset V_1$, $X \subset V'_2 \setminus V_1$, $|X| + |W_1| = k$ and C_1 is a connected component of $G_1|_{W_1}$. Let $\phi_1 : A_1 \rightarrow B_1$ be the map defined by $\phi_1(C, W) = (C \cap V_1, W \cap V_1, W \setminus V_1)$. Then ϕ_1 has the inverse map defined as follows. For a triple $(C_1, W_1, X) \in B_1$, $\phi_1^{-1}(C_1, W_1, X) = (C, W)$, where $W = W_1 \cup X$ and C is the connected component of $G|_W$ containing C_1 . Thus ϕ_1 is a bijection and we get $|A_1| = |B_1| = \sum_{i=0}^k c_i(G_1) \binom{n_2-t}{k-i}$. Similarly we get $|A_2| = \sum_{i=0}^k c_i(G_2) \binom{n_1-t}{k-i}$.

Now let $B = \{W \subset V_1 \cup V'_2 : W \cap F_1 \neq \emptyset\}$. Let $\psi : A_1 \cap A_2 \rightarrow B$ be the map defined by $\psi(C, W) = W$. We have the inverse map ψ^{-1} as follows. For $W \in B$, $\psi^{-1}(W) = (C, W)$, where C is the connected component of $G|_W$ containing $W \cap F_1$, which is guaranteed to exist since $G|_{F_1} = G_1|_{F_1}$ is a complete graph. Thus ψ is a bijection, and we get $|A_1 \cap A_2| = |B| = \binom{n_1+n_2-t}{k} - \binom{n_1+n_2-2t}{k}$. \square

Theorem 3.2. *Let G_1 and G_2 be graphs with n_1 and n_2 vertices respectively. Let t be a positive integer and let G be a t -connected sum of G_1 and G_2 . Then*

$$b_k(G) = \sum_{i=0}^k \left(b_i(G_1) \binom{n_2-t}{k-i} + b_i(G_2) \binom{n_1-t}{k-i} \right) + \binom{n_1+n_2-2t}{k}.$$

Proof. Since $c_k(G) = b_k(G) + \binom{n_1+n_2-t}{k}$, $c_i(G_1) = b_i(G_1) + \binom{n_1}{i}$ and $c_i(G_2) = b_i(G_2) + \binom{n_2}{i}$, by Lemma 3.1, it is sufficient to show that

$$2\binom{n_1+n_2-t}{k} = \sum_{i=0}^k \left(\binom{n_1}{i} \binom{n_2-t}{k-i} + \binom{n_2}{i} \binom{n_1-t}{k-i} \right),$$

which is immediate from the identity $\sum_{i=0}^k \binom{a}{i} \binom{b}{k-i} = \binom{a+b}{k}$. \square

Recall that a t -connected sum G of two graphs depends on the choice of vertices of each graph and the identification of the chosen vertices. However, Theorem 3.2 says that $b_k(G)$ does not depend on them. Thus we get the following important property of a t -connected sum of graphs.

Corollary 3.3. *Let t be a positive integer and let G be a t -connected sum of graphs G_1, G_2, \dots, G_n . If H is also a t -connected sum of G_1, G_2, \dots, G_n , then $b_k(G) = b_k(H)$ for all k .*

Using Proposition 2.3, we get a formula for the special graded Betti number of a t -connected sum of two simplicial complexes for $t \geq 3$.

Corollary 3.4. *Let Δ_1 and Δ_2 be simplicial complexes on V_1 and V_2 respectively with $|V_1| = n_1$ and $|V_2| = n_2$. Let t be a positive integer and let Δ be a t -connected sum of Δ_1 and Δ_2 . If $t \geq 3$ then*

$$b_k(\Delta) = \sum_{i=0}^k \left(b_i(\Delta_1) \binom{n_2-t}{k-i} + b_i(\Delta_2) \binom{n_1-t}{k-i} \right) + \binom{n_1+n_2-2t}{k}.$$

For an integer n , let K_n denote a complete graph with n vertices.

Let G be a graph with vertex set V . If H is a t -connected sum of G and K_{t+1} then H is a graph obtained from G by adding a new vertex v connected to all vertices in W for some $W \subset V$ such that $G|_W$ is isomorphic to K_t . Thus H is determined by choosing such a subset $W \subset V$. Using this observation, we get the following lemma.

Theorem 3.5. *Let t be a positive integer. Let G be a t -connected sum of n K_{t+1} 's. Then*

$$b_k(G) = (k-1) \binom{n}{k}.$$

Proof. We construct a sequence of graphs H_1, \dots, H_n as follows. Let H_1 be the complete graph with vertex set $\{v_1, v_2, \dots, v_{t+1}\}$. For $i \geq 2$, let H_i be the graph obtained from H_{i-1} by adding a new vertex v_{t+i} connected to all vertices in $\{v_1, v_2, \dots, v_t\}$. Then H_n is a t -connected sum of n K_{t+1} 's, and we have $b_k(G) = b_k(H_n)$ by Corollary 3.3. In H_n , the vertex v_i is connected to all the other vertices for $i \leq t$, and v_j and $v_{j'}$ are not connected to each other for all $t+1 \leq j, j' \leq t+n$. Thus $b_k(H_n) = (k-1) \binom{n}{k}$. \square

Observe that every tree with $n+1$ vertices is a 1-connected sum of n K_2 's. Thus we get the following nontrivial property of trees which was observed by Bruns and Hibi [2].

Corollary 3.6. [2, Example 2.1. (b)] *Let T be a tree with $n+1$ vertices. Then $b_k(T)$ does not depend on the specific tree T . We have*

$$b_k(T) = (k-1) \binom{n}{k}.$$

Corollary 3.7. [2, Example 2.1. (c)] *Let G be an n -gon. If $k = n$ then $b_k(G) = 0$; otherwise,*

$$b_k(G) = \frac{n(k-1)}{n-k} \binom{n-2}{k}.$$

Proof. It is clear for $k = n$. Assume $k < n$. Let $V = \{v_1, \dots, v_n\}$ be the vertex set of G . Then

$$\begin{aligned} (n-k) \cdot b_k(G) &= \sum_{\substack{W \subset V \\ |W|=k}} (\text{cc}(G|_W) - 1) \sum_{v \in V \setminus W} 1 \\ &= \sum_{v \in V} \sum_{\substack{W \subset V \setminus \{v\} \\ |W|=k}} (\text{cc}(G|_W) - 1) \\ &= \sum_{v \in V} b_k(G|_{V \setminus \{v\}}). \end{aligned}$$

Since each $G|_{V \setminus \{v\}}$ is a tree with $n-1$ vertices, we are done by Corollary 3.6. \square

Remark 3.8. Bruns and Hibi [2] obtained Corollary 3.6 and Corollary 3.7 by showing that if Δ is a tree (or an n -gon), considered as a 1-dimensional simplicial complex, then $\mathbf{k}[\Delta]$ has a pure resolution. Since $\mathbf{k}[\Delta]$ is Cohen-Macaulay and it has a pure resolution, the Betti numbers are determined by its type (c.f. [1]).

Now we can prove (3). Note that, for $d \geq 3$, if P is a d -dimensional simplicial polytope and Q is a simplicial polytope obtained from P by attaching a d -dimensional simplex S to a facet of P , then $\Delta(Q)$ is a d -connected sum of $\Delta(P)$ and $\Delta(S)$, and thus the 1-skeleton $G(\Delta(Q))$ is a d -connected sum of $G(\Delta(P))$ and K_{d+1} . Hence the 1-skeleton of the boundary complex of a d -dimensional stacked polytope is a d -connected sum of K_{d+1} 's.

Theorem 3.9. *Let P be a d -dimensional stacked polytope with n vertices. If $d \geq 3$, then*

$$b_k(\Delta(P)) = (k-1) \binom{n-d}{k}.$$

If $d = 2$, then

$$b_k(\Delta(P)) = \begin{cases} 0, & \text{if } k = n, \\ \frac{n(k-1)}{n-k} \binom{n-2}{k}, & \text{otherwise.} \end{cases}$$

Proof. Assume $d \geq 3$. Then the 1-skeleton $G(\Delta(P))$ is a d -connected sum of $n-d$ K_{d+1} 's. Thus by Theorem 3.5, we get $b_k(\Delta(P)) = b_k(G(\Delta(P))) = (k-1) \binom{n-d}{k}$.

Now assume $d = 2$. Then $G(\Delta(P))$ is an n -gon. Thus by Corollary 3.7 we are done. \square

REFERENCES

- [1] Winfried Bruns and Jürgen Herzog. *Cohen-Macaulay rings*, volume 39 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1993.
- [2] Winfried Bruns and Takayuki Hibi. Cohen-Macaulay partially ordered sets with pure resolutions. *European J. Combin.*, 19(7):779–785, 1998.
- [3] Victor M. Buchstaber and Taras E. Panov. *Torus actions and their applications in topology and combinatorics*, volume 24 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2002.
- [4] Jürgen Herzog and Enzo Maria Li Marzi. Bounds for the Betti numbers of shellable simplicial complexes and polytopes. In *Commutative algebra and algebraic geometry (Ferrara)*, volume 206 of *Lecture Notes in Pure and Appl. Math.*, pages 157–167. Dekker, New York, 1999.

- [5] Richard P. Stanley. *Combinatorics and commutative algebra*, volume 41 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, second edition, 1996.
- [6] Naoki Terai and Takayuki Hibi. Computation of Betti numbers of monomial ideals associated with stacked polytopes. *Manuscripta Math.*, 92(4):447–453, 1997.
- [7] Günter M. Ziegler. *Lectures on polytopes*, volume 152 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.

DEPARTMENT OF MATHEMATICAL SCIENCES, KAIST, 335 GWAHANGNO, YUSEONG-GU, DAEJEON 305-701,
REPUBLIC OF KOREA

E-mail address: choisy@kaist.ac.kr

DEPARTMENT OF MATHEMATICAL SCIENCES, KAIST, 335 GWAHANGNO, YUSEONG-GU, DAEJEON 305-701,
REPUBLIC OF KOREA

E-mail address: jskim@kaist.ac.kr